

Available online at www.sciencedirect.com

Journal of Functional Analysis 251 (2007) 210–231

**JOURNAL OF
Functional
Analysis**

www.elsevier.com/locate/jfa

On C^* -algebra of a semigroup of partial isometries

Vasyl Ostrovskiy^a, Stanislav Popovych^b, Lyudmila Turowska^{b,*}

^a *Institute of Mathematics, Tereshchenkivska 3, Kyiv 01601, Ukraine*

^b *Mathematical Sciences, Chalmers University of Technology, Mathematical Sciences,
Göteborg University, SE-412 96 Göteborg, Sweden*

Received 13 December 2006; accepted 26 March 2007

Available online 4 May 2007

Communicated by D. Voiculescu

Abstract

A C^* -algebra associated to strongly continuous one-parameter semigroups of partial isometries is introduced as a groupoid C^* -algebra $C^*(\mathcal{G})$. A one-to-one correspondence between non-degenerate representations of $C^*(\mathcal{G})$ and the semigroups is established. It is proved also that $C^*(\mathcal{G})$ contains an ideal J isomorphic to $C_0((0, +\infty], \mathcal{K})$ and $C^*(\mathcal{G})/J \simeq C_0(\mathbb{R})$, where \mathcal{K} is the algebra of compact operators. The C^* -algebra $C^*(\mathcal{G})$ is shown to be generated by one unbounded affiliated element whose image under each non-degenerate representation is the infinitesimal generator of the corresponding semigroup.

© 2007 Elsevier Inc. All rights reserved.

Keywords: Semigroups; Partial isometries; Groupoids; Groupoid C^* -algebra; Representations

1. Introduction

Let $t \mapsto U_t$ for $t \geq 0$ be a strongly continuous one-parameter semigroup of partial isometries on a Hilbert space H . An example of such a semigroup $t \mapsto R_t^a$ which is not unitary not isometric and not coisometric is given by considering for $a > 0$ the Hilbert space $H = L_2([0, a])$ and

$$(R_t^a \xi)(x) = \begin{cases} \xi(x - t), & \text{if } t \leq x \leq a, \\ 0, & \text{if } x < t, \end{cases}$$

* Corresponding author. Fax: +46 31 161973.

E-mail addresses: vo@imath.kiev.ua (V. Ostrovskiy), stanislsp@math.chalmers.se (S. Popovych), turowska@math.chalmers.se (L. Turowska).

with the understanding that $R_t^a = 0$ if $t > a$. Any semigroup $t \mapsto U_t$ which is unitarily equivalent to R_t^a is called a truncated shift.

Strongly continuous one-parameter semigroups of partial isometries were studied in [8,12], where a complete structure theorem was obtained. It asserts, in analogy with the Wold decomposition of single isometry, that for $t \mapsto U_t$, $t \geq 0$, there are Hilbert spaces H_1 , H_+ , H_- , H_0 and there is a unitary operator S such that

$$SU_tS^* = V_t \oplus K_t \oplus S_t \oplus T_t,$$

where V_t is unitary on H_1 , K_t is purely isometric on H_+ , S_t is purely coisometric on H_- and T_t on H_0 is a direct integral of truncated shifts. The structure of more general centered one-parameter semigroups were studied in [12].

Recall that there exists a unique irreducible semigroup of isometries $t \mapsto K_t$ up to unitary equivalence which is given by $(K_t\xi)(x) = \chi_{[t,+\infty)}(x)\xi(x-t)$, $\xi \in L_2([0,+\infty))$. The semigroup is important in the study of Wiener–Hopf equations; i.e., equations of the form

$$(I + W(f))\xi = \eta,$$

where $f \in L_1(\mathbb{R})$, $\eta, \xi \in L_2([0,+\infty))$ and where $W(f)$ is the Wiener–Hopf operator defined by the formula

$$(W(f)\xi)(x) = \int_{\mathbb{R}} f(t)(K_t\xi)(x)dt = \int_0^{+\infty} f(x-t)\xi(t)dt.$$

The Wiener–Hopf C^* -algebra \mathcal{W} generated by all Wiener–Hopf operators is the universal C^* -algebra for semigroups of isometries in the sense that every its non-degenerate representation $\pi : \mathcal{W} \rightarrow B(H)$ arises from unique semigroup of isometries $t \mapsto U_t$: $\pi(W(f)) = \int_{\mathbb{R}} f(t)U_t dt$ [4,5]. This C^* -algebra and some its generalisations were studied intensively in the literature in connection with numerous applications [1,4,5,10]. There is no approach in the literature to associate a C^* -algebra to a general strongly continuous one-parameter semigroup $t \mapsto U_t$, $t \geq 0$, of partial isometries (for the case of discrete semigroups $t \mapsto U_t$ see [9]). The aim of this work is to fill up this gap. We will use a groupoid C^* -algebras approach which was introduced by Muhly and Renault [10] in connection with the study of the C^* -algebras of Wiener–Hopf operators on normal subsemigroups of locally-compact groups.

Consider for $t \mapsto U_t$, $t \geq 0$, the C^* -algebra $\mathfrak{A}(U_t : t \geq 0)$ generated by Wiener–Hopf type operators $\int_{\mathbb{R}} f(t)U_t dt$ ($f \in L_1(\mathbb{R})$, $U_{-t} = U_t^*$, $t > 0$). In this paper we show that for U_t which is a direct integral of R_t^a with respect to the Lebesgue measure on $(0,+\infty)$, $\mathfrak{A}(U_t : t \geq 0)$ is the image of a faithful representation of a groupoid C^* -algebra $C^*(\mathcal{G})$ for a certain locally compact groupoid \mathcal{G} . We establish a one-to-one correspondence between the representations of $C^*(\mathcal{G})$ and strongly continuous one-parameter semigroups of partial isometries up to a natural action of \mathbb{Z}^2 given by the involution: $t \mapsto U_t$, $t \geq 0$, is a semigroup of partial isometries iff so is the family $t \mapsto U_t^*$, $t \geq 0$. To obtain this fact we established the exact sequence

$$0 \rightarrow C_0((0,+\infty], \mathcal{K}) \rightarrow C^*(\mathcal{G}) \rightarrow C_0(\mathbb{R}) \rightarrow 0.$$

Here \mathcal{K} is the algebra of compact operators. Finally we found an unbounded generator A of the C^* -algebra $C^*(\mathcal{G})$ in the sense of [16]. It has the property that for each non-degenerated

representation π of $C^*(\mathcal{G})$ the operator $\pi(A)$ is the generator of the corresponding one-parameter semigroup of partial isometries.

2. Preliminary

For $a > 0$ consider the following strongly continuous one-parameter family of partial isometries acting on $L_2([0, a])$:

$$(U_t^a \xi)(x) = \begin{cases} \chi_{[t, a]}(x) \xi(x - t), & \text{if } t \geq 0, \\ \chi_{[0, a+t]}(x) \xi(x - t), & \text{if } t < 0, \end{cases}$$

where χ_A is the indicator of the set A . One can check that $U_t^a U_s^a = U_{t+s}^a$, $(U_t^a)^* = U_{-t}^a$ for all $t, s \in \mathbb{R}^+$ and that U_t^a is a partial isometry with the initial projection $M_{\chi_{[0, a-t]}}$ and the range projection $M_{\chi_{[t, a]}}$ (for $t > 0$). Here M_g is the multiplication operator on $L_2([0, a])$ by $g \in L_\infty([0, a])$. We also denote by $\{U_t^\infty\}_{t \in \mathbb{R}}$ the strongly continuous family of isometries acting on $L_2([0, +\infty))$ by

$$(U_t^\infty \xi)(x) = \begin{cases} \chi_{[t, +\infty)}(x) \xi(x - t), & \text{if } t \geq 0, \\ \xi(x - t), & \text{if } t < 0. \end{cases}$$

By [8,12] any irreducible strongly continuous one-parameter semigroup of partial isometries, $\{u_t\}_{t \geq 0}$, $u_0 = 1$, on a Hilbert space is unitarily equivalent either to $\{U_t^a\}_{t \geq 0}$, $a \in \mathbb{R}^+ \cup \{\infty\}$ or to $\{(U_t^\infty)^*\}_{t \geq 0} = \{U_{-t}^\infty\}_{t \geq 0}$ or it is a one-dimensional unitary semigroup $\{e^{it\phi}\}_{t \geq 0}$, $\phi \in \mathbb{R}$.

Note that $\{(U_t^a)^*\}_{t \geq 0}$, $0 < a < \infty$, is also an irreducible strongly continuous one-parameter semigroup of partial isometries. It is unitarily equivalent to $\{U_t^a\}_{t \geq 0}$, the equivalence is given by $(W\xi)(x) = \xi(a - x)$.

To each $f \in L_1(\mathbb{R})$ let us associate an operator $W_a(f)$ on $L_2([0, a])$ if $0 < a < \infty$ or on $L_2([0, +\infty))$ if $a = \infty$ given by

$$(W_a(f)\xi)(x) = \int_{\mathbb{R}} f(s) U_s^a \xi(x) ds = \int_{x-a}^x f(s) \xi(x-s) ds$$

($x - \infty := -\infty$ for any $x \in \mathbb{R}$). The operators $W_\infty(f)$ are known as Wiener–Hopf operators. The C^* -algebra generated by all those operators, also called the Wiener–Hopf C^* -algebra, was extensively studied in the literature, in particular, because of its numerous applications, see [1,3–5,7]. In [10] this algebra was realized as the image of a groupoid C^* -algebra under a suitable representation. Our aim is to associate a groupoid C^* -algebra to the semigroup of partial isometries following an idea similar to one in [10]. For this let us calculate first the product $W_a(f_1)W_a(f_2)$, $f_1, f_2 \in L_1(\mathbb{R})$ to find the following:

$$\begin{aligned} (W_a(f_1)W_a(f_2)\xi)(x) &= \int_{x-a}^x f_1(t) \int_{x-t-a}^{x-t} f_2(s) \xi(x-t-s) ds dt \\ &= \int_{x-a}^x \int_{x-a}^x f_1(t) f_2(s-t) \xi(x-s) ds dt \end{aligned}$$

$$\begin{aligned}
&= \int_{x-a}^x \left(\int_{x-a}^x f_1(t) f_2(s-t) dt \right) \xi(x-s) ds \\
&= \int_{x-a}^x h(x, s, a) \xi(x-s) ds,
\end{aligned}$$

where $h(x, s, a) = \int_{x-a}^x f_1(t) f_2(s-t) dt$. Take now $f_i \in L_1(\mathbb{R})$, $i = 1, 2, 3, 4$, and let h_1 and h_2 correspond to $W_a(f_1)W_a(f_2)$ and $W_a(f_3)W_a(f_4)$, respectively. Then we have

$$\begin{aligned}
&(W_a(f_1)W_a(f_2)W_a(f_3)W_a(f_4)\xi)(x) \\
&= \int_{x-a}^x h_1(x, s, a) \int_{x-s-a}^{x-s} h_2(x-s, u, a) \xi(x-s-u) du ds \\
&= \int_{x-a}^x h_1(x, s, a) \int_{x-a}^x h_2(x-s, u-s) \xi(x-u) du ds \\
&= \int_{x-a}^x h(x, u, a) \xi(x-u) du,
\end{aligned}$$

where

$$h(x, u, a) = \int_{x-a}^x h_1(x, s, a) h_2(x-s, u-s, a) ds. \quad (1)$$

Given $a \in (0, \infty]$, let $X_a = [0, a]$ if $a > 0$ and $X_a = [0, +\infty)$ if $a = \infty$. Consider the collection \mathfrak{H} of all functions h on $[0, +\infty) \times \mathbb{R} \times (0, +\infty]$ such that for each $a \in (0, +\infty]$ $\text{supp } h(\cdot, \cdot, a) \subseteq X_a$ and

$$\text{ess sup} \left\{ \int_{x-a}^x h(x, s, a) ds \mid a > 0 \cup a = \infty, x \in [0, a] \right\} \quad (2)$$

is finite. An easy calculation shows that $W_a(h)^* = W_a(h^*)$, where

$$h^*(x, s, a) = \overline{h(x-s, -s, a)}.$$

With respect to the product defined by (1) and expression (2) as norm \mathfrak{H} becomes a $*$ -Banach algebra. The operators

$$(W_a(h)\xi)(x) = \int_{x-a}^x h(x, s, a) \xi(x-s) ds, \quad h \in \mathfrak{H},$$

define a representation of \mathfrak{H} .

3. Groupoids and their C^* -algebras

Let us recall some basic definitions and facts from the theory of groupoids and groupoid C^* -algebras (for detailed treatment see [13,14]).

A *groupoid* is a set \mathfrak{G} together with a subset $\mathfrak{G}^2 \subseteq \mathfrak{G} \times \mathfrak{G}$, a product map $(a, b) \rightarrow ab$ from \mathfrak{G}^2 to \mathfrak{G} and an inverse map $a \rightarrow a^{-1}$ (so that $(a^{-1})^{-1} = a$) such that:

- (i) if $(a, b), (b, c) \in \mathfrak{G}^2$ then $(ab, c), (a, bc) \in \mathfrak{G}^2$ and

$$(ab)c = a(bc).$$

- (ii) $(b, b^{-1}) \in \mathfrak{G}^2$ for all $b \in \mathfrak{G}$ and if $(a, b) \in \mathfrak{G}^2$ then

$$a^{-1}(ab) = b, \quad (ab)b^{-1} = a.$$

If, in addition, \mathfrak{G} is a locally compact Hausdorff space and if the maps defined above are continuous (\mathfrak{G}^2 is endowed with relative topology) then \mathfrak{G} is called a *locally compact groupoid*. The map $d: \mathfrak{G} \rightarrow \mathfrak{G}, x \mapsto x^{-1}x$, is called the *domain map*. The map $r: \mathfrak{G} \rightarrow \mathfrak{G}, x \mapsto xx^{-1}$, is called the *range map*. They have a common image \mathfrak{G}^0 called the *unit space* of \mathfrak{G} .

If \mathfrak{G} is a groupoid and $u \in \mathfrak{G}^0$, then $d^{-1}(u) \cap r^{-1}(u)$ becomes a locally compact group with identity u . This group is called the *isotropy group* at u . Two points u and v in \mathfrak{G}^0 lie in the same orbit if and only if $d^{-1}(u) \cap r^{-1}(v) \neq \emptyset$. For $u \in \mathfrak{G}^0$ the orbit $\text{Orb}(u)$ through u is the set of units that lie in the same orbit with u . A set which is the union of orbits is called *invariant*. A groupoid is called *transitive* if there is only one orbit and *principal* if the map $x \mapsto (d(x), r(x))$ is one-to-one. Let $C_c(\mathfrak{G})$ be the space of compactly supported continuous complex-valued functions defined on a groupoid \mathfrak{G} . A family of measures $\{\lambda^u\}$ on \mathfrak{G} indexed by \mathfrak{G}^0 is called a (left) *Haar system* on \mathfrak{G} if:

- (i) each λ^u is supported by $r^{-1}(u)$,
(ii) for each $f \in C_c(\mathfrak{G})$ the function $u \mapsto \int f d\lambda^u$ is continuous on \mathfrak{G}^0 ,
(iii) for each $x \in \mathfrak{G}$ and $f \in C_c(\mathfrak{G})$, $\int f(xy) d\lambda^{d(x)}(y) = \int f(y) d\lambda^{r(x)}(y)$.

Next we want to associate a groupoid with the algebra \mathfrak{H} from Section 2. Let $\overline{\mathbb{R}}$ denote $(-\infty, \infty]$. Consider the space $\mathcal{G} = \{(x, s, a) \in [0, \infty] \times \mathbb{R} \times (0, \infty]: x \in [0, a], x - s \in [0, a]\}$ with the usual topology on it.

The space becomes a locally compact groupoid if we set $\mathcal{G}^2 = \{[(x, s, a), (y, t, b)] \mid y = x - s, a = b\}$ and define the product and the inverse maps by

$$(x, s, a) \cdot (x - s, t, a) = (x, s + t, a), \quad (x, s, a)^{-1} = (x - s, -s, a).$$

(We use the convention $\infty - s = \infty$ for each $s \in \mathbb{R}$.) The maps d and r satisfy the equations

$$d(x, s, a) = (x, s, a)^{-1}(x, s, a) = (x - s, 0, a), \quad (3)$$

$$r(x, s, a) = (x, s, a)(x, s, a)^{-1} = (x, 0, a). \quad (4)$$

The unit space of \mathcal{G} is $\{(x, 0, a): x \in [0, a], a > 0\} \cup \{(x, 0, \infty): x \geq 0\} \cup \{\infty, 0, \infty\}$.

Let $C_c(\mathcal{G})$ be the space of compactly supported, continuous, complex-valued functions defined on \mathcal{G} . We may take $\lambda^u = \chi_{\mathcal{G}} \delta_x \times \lambda \times \delta_a$ for $u = (x, 0, a)$, where δ_x is a point mass at x and λ is the Lebesgue measure on \mathbb{R} , and obtain a left Haar system on \mathcal{G} . A multiplication, involution and a norm with respect to which $C_c(\mathcal{G})$ becomes a normed $*$ -algebra are defined as follows: for $u = (x, s, a)$, with $a, x \neq \infty$, $f, g \in C_c(\mathcal{G})$ the product is given by

$$\begin{aligned} (f * g)(u) &= \int f(uv)g(v^{-1})d\lambda^{d(u)}(v) = \int_{x-s-a}^{x-s} f(x, s+t, a)g(x-s-t, -t, a)dt \\ &= \int_{x-a}^x f(x, r, a)g(x-r, s-r, a)dr, \end{aligned}$$

otherwise

$$(f * g)(u) = \begin{cases} \int_{-\infty}^x f(x, r, \infty)g(x-r, s-r, \infty)dr, & u = (x, s, \infty), 0 \leq x < \infty, \\ \int_{-\infty}^{+\infty} f(\infty, r, \infty)g(\infty, s-r, \infty)dr, & u = (\infty, s, \infty); \end{cases}$$

the involution and the norm are given by the formulas

$$f^*(u) = \overline{f(u^{-1})} = \overline{f(x-s, -s, a)}$$

and

$$\begin{aligned} \|f\|_I &= \max \left\{ \sup_{u \in \mathcal{G}^0} \int |f| d\lambda^u, \sup_{u \in \mathcal{G}^0} \int |f^*| d\lambda^u \right\} \\ &= \max \left\{ \sup \left\{ \int_{x-a}^x |f(x, t, a)| dt, \int_{x-a}^x |f(x-t, -t, a)| dt : (x, a) \in [0, a] \times (0, \infty) \right\} \right\}. \end{aligned}$$

Let $L^I(\mathcal{G}, \lambda)$ be the completion of $C_c(\mathcal{G})$ with respect to the defined norm. We denote by $C^*(\mathcal{G})$ (or $C^*(\mathcal{G}, \lambda)$) its enveloping C^* -algebra.

4. Structure and representations of the groupoid C^* -algebra $C^*(\mathcal{G})$

The aim of this section is to show that there is a one-to-one correspondence between representations of the defined groupoid C^* -algebra and strongly continuous semigroups of partial isometries up to the action of \mathbb{Z}_2 defined by involution: if $\{u_t\}_{t \geq 0}$ is a strongly continuous semigroup of partial isometries so is the family $\{u_t^*\}_{t \geq 0}$.

In what follows we denote by $C_c(X)$ the algebra of all continuous complex-valued functions on a locally compact set X with compact support and by $C_0(X)$ the continuous complex-valued functions vanishing at infinity.

For $g \in C_c(\mathbb{R} \times (0, +\infty])$ we define $\tilde{g} \in C_c(\mathcal{G})$ by $\tilde{g}(x, s, a) = g(s, a)$.

Proposition 1. *The subalgebra \mathcal{A} generated by $\{\tilde{g} : g \in C_c(\mathbb{R} \times (0, \infty))\}$ is dense in $C^*(\mathcal{G})$.*

Proof. It is enough to prove that \mathcal{A} is dense in $C_c(\mathcal{G})$ endowed with the inductive limit topology. Let $D = [0, \infty] \times (0, \infty]$. We claim first that the algebra generated by $\psi_f(y, a) = \int_{y-a}^y f(t) dt$, $f \in C_c(\mathbb{R})$ is dense in $C_0(D)$. The family of functions $\{\psi_f\}_{f \in C_c(\mathbb{R})}$ separates the points of D . Indeed if $(y, a), (x, b) \in D$ and $\int_{y-a}^y f(t) dt = \int_{x-b}^x f(t) dt$ for all functions $f \in C_c(\mathbb{R})$ then $[y-a, y] = [x-b, x]$ and therefore $(y, a) = (x, b)$. The family $\{\psi_f\}$ does not vanish at any point of $(y, a) \in D$ since otherwise $\int_{y-a}^y f(t) dt = 0$ for all $f \in C_c(\mathbb{R})$ which is impossible if $a > 0$. Clearly, the family is closed under complex conjugation. Thus, by the Stone–Weierstrass theorem the algebra generated by $\{\psi_f\}$ is dense in $C_0(D)$. Hence to prove the statement it is sufficient to approximate the functions of the form $\varphi(x, s, a) = (\int_{x-a}^x f(t) dt)g(s)$, where $f \in C_c(\mathbb{R})$, $g \in C_c(\mathbb{R} \times (0, \infty])$, by the elements of \mathcal{A} . Fix now $f \in C_c(\mathbb{R})$, $g \in C_c(\mathbb{R} \times (0, \infty])$ and denote by K a compact neighborhood of $(\text{supp } g)_1$, here $(\text{supp } g)_1 = \{s: (s, a) \in \text{supp } g \text{ for some } a \in (0, \infty]\}$. Fix $\varepsilon > 0$ and let $M = \int_{\mathbb{R}} |f(t)| dt$ which is finite since f has a compact support. Then there exists a neighborhood $U \subset \mathbb{R}$ of 0 such that $U + \text{supp}(g)_1 \subset K$ and $|g(t, a) - g(s, a)| < \varepsilon/M$ whenever $t - s \in U$. Since $\text{supp}(f)$ is compact, there exists $\{t_i\}_{i=1}^n$, such that $\{U + t_i\}_{i=1}^n$ is an open covering of $\text{supp}(f)$. Let $\{h_i\}_{i=1}^n$ be a partition of unity of $\text{supp}(f)$ such that $\text{supp}(h_i) \subset U + t_i$. We have $\sum_{i=1}^n h_i(r) = 1$ for $r \in \text{supp}(f)$. The set $(\text{supp } g)_2 = \{a: (s, a) \in \text{supp } g \text{ for some } s \in \mathbb{R}\}$ is compact. Let θ be a function in $C_c(\mathbb{R})$ such that $\theta(a) = 1$ for $a \in (\text{supp } g)_2$. Set

$$f_i(r, a) = f(r)\theta(a)h_i(r), \quad g_i(r, a) = g(r + t_i, a)$$

and

$$\psi = \sum_{i=1}^n \tilde{f}_i * \tilde{g}_i.$$

We have

$$\begin{aligned} \varphi(x, s, a) - \psi(x, s, a) &= \left(\int_{x-a}^x f(v) dv \right) g(s, a) - \left(\sum_{i=1}^n \tilde{f}_i * \tilde{g}_i \right)(x, s, a) \\ &= \int_{x-a}^x \left[f(v)g(s, a) - \sum_{i=1}^n \tilde{f}_i(x, v, a) \tilde{g}_i(x - v, s - v, a) \right] dv \\ &= \int_{x-a}^x \left[f(v)g(s, a) - \sum_{i=1}^n f_i(v)g_i(s - v, a) \right] dv \\ &= \sum_{i=1}^n \int_{x-a}^x f_i(v, a)(g(s, a) - g_i(s - v, a)) dv \\ &= \sum_{i=1}^n \int_{x-a}^x f(v)h_i(v)(g(s, a) - g(t_i - v + s, a)) dv. \end{aligned}$$

Moreover, $|g(s, a) - g(t_i - v + s, a)| < \varepsilon/M$ as $s - (t_i - v + s) = v - t_i \in U$ for $v \in \text{supp}(f_i) \subset \text{supp}(h_i) \subset t_i + U$. Thus

$$|\varphi(x, s, a) - \psi(x, s, a)| < (\varepsilon/M) \int_{-a}^a |f(v)| dv < \varepsilon.$$

It is left to see that ψ and φ have common compact support. Clearly, φ is supported in $[0, \infty] \times K \times (\text{supp } g)_2$. Since $f h_i$ is supported in $U + t_i$ then the integrand $f_i(v, a) g_i(s - v, a) = f(v) h_i(v) g(t_i + s - v, a)$ vanishes whenever $v \notin U + t_i$. Further $t_i + s - v \in (\text{supp } g)_1$ if and only if $s \in v - t_i + (\text{supp } g)_1$. Therefore if $v \in U + t_i$ and $t_i + s - v \in (\text{supp } g)_1$ then $s \in U + t_i - t_i + (\text{supp } g)_1 \subset U + (\text{supp } g)_1 \subset K$. Thus, ψ is also supported in $[0, \infty] \times K \times (\text{supp } g)_2$. This completes the proof. \square

For the chosen left Haar system $\{\lambda^u\}$ on \mathcal{G} let $\lambda_u = (\lambda^u)^{-1}$ be the image of λ^u under the map $x \rightarrow x^{-1}$. Then every Radon measure μ on \mathcal{G}^0 induces two measures ν and ν^{-1} on \mathcal{G} by the rules $\nu = \int \lambda^u d\mu(u)$, $\nu^{-1} = \int \lambda_u d\mu(u)$.

The Hilbert space $L^2(\nu^{-1})$ carries a representation of $C^*(\mathcal{G})$ which is called the representation induced off the unit space by μ and is denoted by $\text{Ind } \mu$. It is defined by the formula

$$\text{Ind } \mu(f) \xi(x) = \int f(xy) \xi(y^{-1}) d\lambda^{d(x)}(y),$$

where $f \in C_c(\mathcal{G})$, $\xi \in L^2(\nu^{-1})$. Since $\|\text{Ind } \mu(f)\| \leq \|f\|_{L^1}$ the representation $\text{Ind } \mu$ extends to $C^*(\mathcal{G})$.

Let $u = (0, 0, a) \in \mathcal{G}^0$ and let δ_u be the point mass at $\{u\}$. Denote the induced measures by ν_u and ν_u^{-1} . Then for $f \in C_c(\mathcal{G})$ we have

$$\int f d\nu_u = \int_{-a}^0 f(0, s, a) ds \quad \text{and} \quad \int f d\nu_u^{-1} = \int_0^a f(s, s, a) ds,$$

giving $\text{supp } \nu_u^{-1} = \{(s, s, a) : s \in \mathbb{R}\}$. For $\xi \in L^2(\nu_u^{-1})$ and $f \in C_c(\mathcal{G})$ we have

$$(\text{Ind } \delta_u(f) \xi)(s, s, a) = \int_0^a f(s, s - t, a) \xi(t, t, a) dt. \quad (5)$$

The operator $U : L^2([0, a]) \rightarrow L^2(\nu_u^{-1})$ defined by the formula

$$(U\xi)(x, t, b) = \begin{cases} \xi(t), & x = t, b = a, \\ 0, & \text{otherwise,} \end{cases}$$

where $\xi \in L^2([0, a], \lambda)$ is an isometry with the inverse $(U^{-1}\zeta)(t) = \zeta(t, t, a)$ ($\zeta \in L^2(\nu_u^{-1})$). For $f \in C_c(\mathcal{G})$ and $\xi \in L^2(\nu_u^{-1})$ we have

$$\begin{aligned}
(UW_a(f)U^{-1}\xi)(t, t, a) &= (W_a(f)U^{-1}\xi)(t) \\
&= \int_{t-a}^t f(t, s, a)(U^{-1}\xi)(t-s) ds \\
&= \int_{t-a}^t f(t, s, a)\xi(t-s, t-s, a) ds \\
&= \int_0^a f(t, t-s, a)\xi(s, s, a) ds.
\end{aligned}$$

Comparing with (5) we obtain that $\text{Ind } \delta_u$ is unitarily equivalent to W_a . Moreover, by [10, Proof of Proposition 2.17], if $u, v \in \mathcal{G}_0$ lie in the same orbit then $\text{Ind } \delta_u$ and $\text{Ind } \delta_v$ are unitarily equivalent, so that for any $a \in (0, \infty]$ and $x, y \neq \infty$, $\text{Ind } \delta_{(x,0,a)} \simeq \text{Ind } \delta_{(y,0,a)}$. That W_a , $a \in (0, \infty]$, is irreducible can be easily checked.

If $u = (\infty, 0, \infty)$, then for $f \in C_c(\mathcal{G})$, $\int f dv_u^{-1} = \int_{\mathbb{R}} f(\infty, s, \infty) ds$ and for $\xi \in L_2(v_u^{-1})$ we have

$$(\text{Ind } \delta_u(f)\xi)(\infty, s, \infty) = \int_{\mathbb{R}} f(\infty, s-t, \infty)\xi(\infty, t, \infty) dt = (\tilde{f} * \tilde{\xi})(s),$$

where $\tilde{f}(s) = f(\infty, t, \infty)$, and $*$ is the usual convolution.

Consider also the following family of one-dimensional representations ω_ϕ , $\phi \in \mathbb{R}$, of $C^*(\mathcal{G})$: for $f \in C_c(\mathcal{G})$,

$$\omega_\phi(f) = \int_{\mathbb{R}} f(\infty, t, \infty) e^{-it\phi} dt = \widehat{\tilde{f}}(\phi),$$

where $\widehat{\tilde{f}}$ is the Fourier transform of \tilde{f} . Those representations come from one-dimensional strongly continuous one-parameter semigroups of partial isometries = unitaries. We have that $\text{Ind } \delta_{(\infty,0,\infty)}$ is unitarily equivalent to the direct integral $\int_{\mathbb{R}}^{\oplus} \omega_\phi d\phi$ of these one-dimensional representations via the Fourier transform on \mathbb{R} .

Note also that by [10, Proposition 2.13], $\text{Ind } \mu \simeq \int_{\mathcal{G}^0}^{\oplus} \text{Ind } \delta_u d\mu$.

Let $C_{\text{red}}^*(\mathcal{G}, \lambda)$ be the reduced groupoid C^* -algebra of \mathcal{G} . Recall that it is obtained by the completion of $C_c(\mathcal{G}, \lambda)$ with respect to the norm $\|f\|_{\text{red}} = \sup \|L(f)\|$, $f \in C_c(\mathcal{G})$, where L ranges over all representations induced off the unit space [14, Chapter II, Definition 2.8]. By [10, Proposition 2.15], $C_{\text{red}}^*(\mathcal{G}, \lambda) \cong C^*(\mathcal{G}, \lambda)$.

Proposition 2. *Let m be a probability measure on \mathbb{R}^+ which is equivalent to the Lebesgue measure. Then the representation $\int_{a>0}^{\oplus} \text{Ind } \delta_{(0,0,a)} dm(a)$ of $C^*(\mathcal{G})$ is faithful.*

Proof. Let λ be our Haar system for \mathcal{G} . By [10, Proposition, p. 12], for a measure μ on \mathcal{G}^0 the kernel of $\text{Ind } \mu = \int_{\mathcal{G}^0}^{\oplus} \text{Ind } \delta_u d\mu(u)$ in $C_{\text{red}}^*(\mathcal{G}, \lambda) = C^*(\mathcal{G}, \lambda)$ is the ideal $C_{\text{red}}^*(\mathcal{G}|_U, 1_U\lambda)$, where U is the largest open invariant subset of \mathcal{G}^0 such that $\mu(U) = 0$. So $\text{Ind } \mu$ has the trivial

kernel if for any open invariant subset U of \mathcal{G}^0 we have $\mu(U) \neq 0$. As $\mathcal{G}^0 = \{(x, 0, a): x \in [0, a], a > 0\} \cup \{(x, 0, \infty): x \geq 0\} \cup \{(\infty, 0, \infty)\}$ and $\text{Orb}((x, 0, a)) = \{(y, 0, a): y \in [0, a]\}$ if $a \in (0, \infty]$, $x \neq \infty$, and $\text{Orb}((\infty, 0, \infty)) = \{(\infty, 0, \infty)\}$, $\ker \text{Ind } \mu = \{0\}$ for $\mu = \delta_0 \times \delta_0 \times m$ as the projection of any open invariant subset U on the third coordinate contains always an interval. \square

Let \mathcal{K} be the C^* -algebra of compact operators acting on a separable Hilbert space. Denote by $C_b(\mathbb{R}^+, \mathcal{K})$ the C^* -algebra of bounded complex-valued continuous functions $f: \mathbb{R}^+ \rightarrow \mathcal{K}$.

Theorem 3. *There exists an embedding of $C^*(\mathcal{G})$ into $C_b((0, +\infty), \mathcal{K})$ and hence $C^*(\mathcal{G})$ is postliminal.*

Proof. We will construct a homomorphism from the subalgebra \mathcal{A} from Proposition 1 to $C_b((0, +\infty), \mathcal{K})$. Let $f \in C_c(\mathbb{R} \times (0, \infty])$. Then $W_a(\tilde{f}) = \int_{-\infty}^{+\infty} f(t, a) U_t^a dt$, where $(U_t^a \xi)(x) = \chi_{[t, a]}(x) \xi(x - t)$ for $t > 0$, $U_t^a = (U_{-t}^a)^*$ for $t < 0$ and $U_0^a = I$. We have shown that each representation W_a is unitarily equivalent to $\text{Ind } \delta_{(0, 0, a)}$. Moreover, $W_a(\tilde{f})$ is compact. Indeed,

$$(W_a(\tilde{f})\xi)(x) = \int_{x-a}^x f(t, a) \xi(x-t) dt = \int_0^a f(x-s, a) \xi(s) ds, \quad \xi \in L_2([0, a]),$$

which is clearly a Hilbert–Schmidt operator.

Define $V_a: L_2([0, 1]) \rightarrow L_2([0, a])$ as follows:

$$(V_a \xi)(x) = \xi\left(\frac{x}{a}\right).$$

Then

$$\|V_a \xi\|^2 = \int_0^a \left| \xi\left(\frac{x}{a}\right) \right|^2 dx = a \int_0^1 |\xi(y)|^2 dy = a \|\xi\|^2$$

and $S_a = \frac{1}{\sqrt{a}} V_a$ is a unitary operator with $(S_a^* \eta)(x) = \sqrt{a} \eta(xa)$, $\eta \in L_2([0, a])$. Moreover,

$$\begin{aligned} (V_t^a \xi)(x) &:= (S_a^* U_t^a S_a \xi)(x) = U_t^a V_a \xi(ax) = \chi_{[t, a]}(ax) (V_a \xi)(ax - t) \\ &= \chi_{[t/a, 1]}(ax) \xi((ax - t)/a) = \chi_{[t/a, 1]}(x) \xi(x - t/a) = (U_{t/a}^1 \xi)(x). \end{aligned}$$

Define now for each $a > 0$ a compact operator $F(a) = S_a^* W_a(\tilde{f}) S_a$. Then $F: (0, \infty) \rightarrow \mathcal{K}$ is continuous and $F(a) \rightarrow 0$ as $a \rightarrow 0$. In fact,

$$S_a^* W_a(\tilde{f}) S_a = \int_{-\infty}^{+\infty} f(t, a) V_t^a dt = \int_{-\infty}^{+\infty} f(t, a) U_{t/a}^1 dt = a \int_{-\infty}^{+\infty} f(as, a) U_s^1 ds$$

and

$$\begin{aligned}\|F(a) - F(b)\| &= \left\| a \int_{-\infty}^{+\infty} f(as, a) U_s^1 ds - b \int_{-\infty}^{+\infty} f(bs, b) U_s^1 ds \right\| \\ &\leq \int_{-1}^1 |af(as, a) - bf(bs, b)| ds.\end{aligned}$$

As f is continuous and compactly supported, $\|F(a) - F(b)\| \rightarrow 0$ as $a \rightarrow b$.

$$\|F(a)\| \leq \int_{-1}^1 |af(as, a)| ds = \int_{-a}^a |f(s, a)| ds \rightarrow 0 \quad \text{as } a \rightarrow 0.$$

Moreover,

$$F(a) \cdot G(a) = S_a^* W_a(\tilde{f}) S_a \cdot S_a^* W_a(\tilde{g}) S_a = S_a^* W_a(\tilde{f} * \tilde{g}) S_a.$$

Define a homomorphism ψ from the algebra generated by \tilde{f} , $f \in C_c(\mathbb{R} \times (0, \infty])$, to $C_0((0, \infty), \mathcal{K})$ by

$$\psi(g)(a) = S_a^* W_a(g) S_a.$$

Then $\|\psi(g)\| = \sup_a \|\psi(g)(a)\| = \sup_a \|W_a(g)\| \leq \|g\|$. Thus ψ can be extended to the whole C^* -algebra $C^*(\mathcal{G})$ by continuity. We have that ψ is injective since $W_a(f) = 0$ for any a if and only if $f = 0$ as the representation $\int_{a>0}^{\oplus} \text{Ind } \delta_{(0,0,a)} dm$ is faithful by Proposition 2. \square

Our aim now will be to show that all the representations of $C^*(\mathcal{G})$ come from the semigroups of partial isometries.

Let $U = \mathcal{G}^0 \setminus \{(\infty, 0, \infty)\}$, which is an open invariant subset in \mathcal{G}^0 . Let J be the closure of $C_c(\mathcal{G}|_U) = \{f \in C_c(\mathcal{G}) \setminus \{(\infty, s, \infty) : s \in \mathbb{R}\}\}$. We have $C_c(\mathcal{G}|_U) \subset \{f \in C_c(\mathcal{G}) : (\text{supp } f)_1 \subset [0, C] \text{ for some } C > 0\}$. By [14, Proposition 4.4], J is a two-sided ideal of $C^*(\mathcal{G})$ and $C^*(\mathcal{G})/J \simeq C^*(\mathcal{G}|_F)$, where $F = \mathcal{G}^0 \setminus U$. Clearly $\mathcal{G}|_F \simeq \mathbb{R}$ so that $C^*(\mathcal{G}|_F) \simeq C^*(\mathbb{R}) \simeq C_0(\mathbb{R})$.

Theorem 4. *The ideal J is isomorphic to $C_0((0, \infty]) \otimes \mathcal{K}$ and $C^*(\mathcal{G})/J \simeq C_0(\mathbb{R})$.*

Thus we have an exact sequence

$$0 \rightarrow C_0((0, \infty]) \otimes \mathcal{K} \rightarrow C^*(\mathcal{G}) \rightarrow C_0(\mathbb{R}) \rightarrow 0.$$

Proof. Let H_a be the Hilbert space $L_2([0, a])$ which we consider as a subspace of $L_2([0, \infty))$ and let

$$\mathcal{B} = \{f \in C_0((0, \infty]) \otimes \mathcal{K} : f(a)H_a \subset H_a, f(a)|_{H_a^\perp} = 0\}.$$

By Proposition A.1, $\mathcal{B} \simeq C_0((0, \infty]) \otimes \mathcal{K}$. Thus it suffices to construct an isomorphism between J and the algebra \mathcal{B} .

For $f \in C_c(\mathcal{G}|_U)$ let $\psi(f)(a) = \widetilde{W}_a(f)$, where

$$(\widetilde{W}_a(f)\xi)(x) = \chi_{[0,a]}(x) \int_{x-a}^x f(x, t, a) \xi(x-t) dt = \int_0^a \chi_{[0,a]}(x) f(x, x-t, a) \xi(t) dt,$$

$\xi \in L_2([0, \infty))$. If P_a denotes the projection from $L_2([0, \infty))$ onto $L_2([0, a])$, then $\widetilde{W}_a(f) = W_a(f)P_a$. If a is finite then clearly $\widetilde{W}_a(f)$ is a Hilbert–Schmidt and therefore compact operator such that $\widetilde{W}_a(f)H_a \subset H_a$ and $\widetilde{W}_a(f)|_{H_a^\perp} = 0$.

Our aim now is to prove that ψ can be extended to an isomorphism $\psi: J \rightarrow \mathcal{B}$. For this we show first that $\psi(f)$ is continuous whenever $f \in J$. We start with proving the convergence $\widetilde{W}_a(f) \rightarrow \widetilde{W}_\infty(f)$ as $a \rightarrow \infty$. Assume that $(\text{supp } f)_1 \subset [0, C]$, $(\text{supp } f)_2 \subset [-A, A]$, $A, C > 0$, and let $a > A + C$. Then

$$\begin{aligned} \|\widetilde{W}_a(f)\xi - \widetilde{W}_\infty(f)\xi\| &= \left(\int_0^{+\infty} \left| \int_0^a (f(x, x-t, \infty) - \chi_{[0,a]}(x) f(x, x-t, a)) \xi(t) dt \right|^2 dx \right)^{1/2} \\ &\quad + \left(\int_a^{+\infty} |f(x, x-t, \infty)|^2 dx \right)^{1/2} \leq I_1 + I_2, \end{aligned}$$

where

$$I_1 = \left(\int_0^{+\infty} \left| \int_0^a (f(x, x-t, \infty) - \chi_{[0,a]}(x) f(x, x-t, a)) \xi(t) dt \right|^2 dx \right)^{1/2}$$

and

$$I_2 = \left(\int_0^{+\infty} \left| \int_a^{+\infty} f(x, x-t, \infty) dt \right|^2 dx \right)^{1/2}.$$

As $(\text{supp } f)_1 \subset [0, C] \subset [0, a - A]$, we have $f(x, x-t, \infty) = 0$ if $x > a - A$. On the other hand, $(\text{supp } f)_2 \subset [-A, A]$ so that $\chi_{[a, +\infty)}(t) f(x, x-t, \infty) = 0$ if $x-t < -A$ and in particular if $x < a - A$. Thus $I_2 = 0$. Using the above condition on the support of f we obtain also that

$$I_1^2 \leq \left(\int_0^C \int_0^{C+A} |f(x, x-t, \infty) - f(x, x-t, a)|^2 dt dx \right) \|\xi\|^2.$$

Thus $\|\widetilde{W}_a(f) - \widetilde{W}_\infty(f)\|^2 \leq \int_0^C \int_0^{C+A} |f(x, x-t, \infty) - f(x, x-t, a)|^2 dt dx$ and since $f(x, x-t, a) \rightarrow f(x, x-t, \infty)$, as $a \rightarrow \infty$, uniformly with respect to (x, t) on each compact set, we obtain $\|\widetilde{W}_a(f) - \widetilde{W}_\infty(f)\| \rightarrow 0$ as $a \rightarrow \infty$. This gives the continuity of $\psi(f)$ at $a = \infty$ and compactness of $\psi(f)(\infty)$. In a similar way one proves the continuity of $\psi(f)$ at $a < \infty$. As $\|\widetilde{W}_a(f)\| = \|W_a(f)\|$ which tends to 0 as $a \rightarrow 0$ (see the proof of Theorem 3) we have now that $\psi(f) \in \mathcal{B}$.

Clearly, ψ is a $*$ -homomorphism from $C_c(\mathcal{G}|_U)$ to \mathcal{B} and $\|\psi(f)(a)\| \leq \|f\|_{C^*(\mathcal{G})}$ for each $f \in C_c(\mathcal{G}|_U)$ and $a \in (0, \infty]$. Thus ψ can be extended to a $*$ -homomorphism from $C^*(\mathcal{G}|_U)$ to \mathcal{B} denoted by the same letter. We have that ψ is injective since $\widetilde{W}_a(f) = 0$ for all a if and only if $f = 0$. The latter follows from the faithfulness of the representation $\int_{a>0}^{\oplus} \text{Ind } \delta_{(0,0,a)} dm$ and the unitary equivalence of W_a and $\text{Ind } \delta_{(0,0,a)}$ for all $a > 0$.

It is left to show now that ψ is surjective. As \mathcal{B} is postliminal it is enough to prove that $\psi(C^*(\mathcal{G}|_U))$ is a rich subalgebra of \mathcal{B} [6, 11.1]. Any irreducible representation of \mathcal{B} is unitarily equivalent to $\pi'_t = \pi_t|_{V_t}$, where $\pi_t(f) = f(t)$, $t \in (0, \infty]$ (see the proof of Proposition A.1). Then $\pi'_t(\psi(f)) = W_t(f)$ for all $f \in C^*(\mathcal{G}|_U)$. Since $\pi'_t \circ \psi|_{C^*(\mathcal{G}|_U)} = W_t|_{C^*(\mathcal{G}|_U)}$ is an irreducible representation of $C^*(\mathcal{G}|_U)$ as the restriction of the irreducible representation W_t of $C^*(\mathcal{G})$ we obtain that $\pi'_t|_{\psi(C^*(\mathcal{G}|_U))}$ is irreducible. Further if π_1, π_2 are non-equivalent irreducible representations of \mathcal{B} then $\pi_1 = \pi'_t, \pi_2 = \pi'_s$, for $t \neq s$ up to a unitary equivalence. As for different t , W_t are non-equivalent representations of $C^*(\mathcal{G})$ and hence of the ideal $C^*(\mathcal{G}|_U)$ we obtain that $\pi_1|_{\psi(C^*(\mathcal{G}|_U))}$ and $\pi_2|_{\psi(C^*(\mathcal{G}|_U))}$ are non-equivalent. Thus $\psi(C^*(\mathcal{G}|_U))$ is a rich subalgebra of \mathcal{B} and the proof is complete. \square

As a corollary we get the following.

Theorem 5. Any irreducible representation of $C^*(\mathcal{G})$ is unitarily equivalent to some W_a , $a \in (0, +\infty]$, or ω_ϕ , $\phi \in \mathbb{R}$.

Proof. It follows from the proof of Theorem 4 that any irreducible representation of J is unitarily equivalent to some $W_a|_J$, $a \in (0, +\infty]$. Hence any irreducible representation of $C^*(\mathcal{G})$ is either unitarily equivalent to W_a , $a \in (0, +\infty]$ or to an irreducible representation which is a composition of an irreducible representation of the quotient algebra $C^*(\mathcal{G})/J \simeq C_0(\mathbb{R})$ and the canonical homomorphism from $C^*(\mathcal{G})$ to $C^*(\mathcal{G})/J$ and thus to a one-dimensional representation ω_ϕ , $\phi \in \mathbb{R}$. \square

Theorem 6. Any representation of $C^*(\mathcal{G})$ is unitarily equivalent to $\bigoplus_{n=1}^{\infty} \text{Ind } \mu_n \otimes I_n \oplus \int_{\mathbb{R}} \omega_\phi \otimes I_n dv_n$ for some Radon measures μ_n on \mathcal{G}^0 and v_n on \mathbb{R} .

Proof. By Theorem 5 the spectrum of $\mathcal{A} = C^*(\mathcal{G})$ as a set is equal to $\{W_a\}_{a \in (0, \infty]} \cup \{\omega_\phi\}_{\phi \in \mathbb{R}}$. It is homeomorphic to a topological space $X = (0, \infty] \sqcup_{\gamma} \mathbb{R}$: here X is a disjoint union of $(0, \infty]$ and \mathbb{R} , $\gamma: \mathbb{R} \rightarrow (0, \infty]$ is a map given by the rule $\gamma(x) = \infty$ for all $x \in \mathbb{R}$, open sets of the topology on X are the open sets U in \mathbb{R} and the sets $V \sqcup \gamma^{-1}(V)$, where V runs over open sets in $(0, \infty]$. The σ -algebra of Borel sets of X is the same as the σ -algebra of Borel sets of the disjoint union $(0, \infty] \sqcup \mathbb{R}$.

Since by Theorem 2, $C^*(\mathcal{G})$ is a postliminal C^* -algebra, any its representation π has a unique decomposition in a direct sum of representations of a fixed multiplicity n , i.e. $\pi = \bigoplus_{n \in \mathbb{N}} \pi_n$. Whereas each π_n is a sum of n copies of multiplicity free representation $\pi^{(n)}$ which is of the form $\int_{\mathcal{A}}^{\oplus} \zeta d\mu^{(n)}(\zeta)$ for some Radon measure $\mu^{(n)}$. In our case we can rewrite the last integral as

$$\int_{(0, \infty]}^{\oplus} W_a d\mu^{(n)}(a) \oplus \int_{\mathbb{R}}^{\oplus} \omega_\phi dv^{(n)}(\phi).$$

If we let μ_n be the measure on \mathcal{G}^0 induced from $\mu^{(n)}$ via the map $\rho: \mathcal{G}^0 \rightarrow (0, \infty]$, $(x, 0, a) \mapsto a$ and let $\nu_n = \nu^{(n)}$ then we obtain the desired decomposition. \square

Let $\text{Rep}(U, H)$ denote the category of strongly-continuous one-parameter semigroups of partial isometries acting on a Hilbert space H and $\text{Rep}(C^*(\mathcal{G}), H)$ is the category of nondegenerate representations of $C^*(\mathcal{G})$ on H .

If $\{U_t\}_{t \geq 0} \in \text{Rep}(U, H)$, then by [8] there exists a unitary S in $B(H)$ such that

$$SU_t S^* = \int_{(0, \infty)}^{\oplus} U_t^{(a)} \otimes I_a d\mu(a) \oplus U_t^\infty \otimes I_\infty \oplus U_t^{\infty*} \otimes I_{-\infty} \oplus \int_{\mathbb{R}}^{\oplus} e^{it\lambda} \otimes I_\lambda d\nu(\lambda), \quad (6)$$

here I_a, I_λ, I_∞ and $I_{-\infty}$ are the identity operators acting on some Hilbert spaces H_a, H_λ, H_∞ and $H_{-\infty}$. We permit H_∞ and $H_{-\infty}$ to be zero-dimensional; in this case the corresponding summand $U_t^\infty \otimes I_\infty$ or $U_t^{\infty*} \otimes I_{-\infty}$ will be omitted. Let $\text{Rep}_+(U, H) = \text{Rep}(U, H)/\mathbb{Z}_2$. The elements of $\text{Rep}_+(U, H)$ can be identified with those semigroups $\{U_t\}_{t \geq 0}$ for which decomposition (6) holds without the components $U_t^{\infty*} \otimes I_{-\infty}$. For such $\{U_t\}_{t \geq 0} \in \text{Rep}_+(U, H)$ we associate a representation $F(\{U_t\})$ of $C^*(\mathcal{G})$ of the form

$$S^* \int_{(0, \infty)}^{\oplus} W_a \otimes I_a d\mu(a) \oplus W_\infty \otimes I_\infty \oplus \int_{\mathbb{R}}^{\oplus} \omega_\lambda \otimes I_\lambda d\nu(\lambda) S.$$

Corollary 1. *The mapping $F: \text{Rep}_+(U, H) \rightarrow \text{Rep}(C^*(\mathcal{G}), H)$ is well defined.*

This follows from the fact that operator S in decomposition (6) of U_t is defined uniquely up to multiplication by a diagonal unitary operator

$$V = \int_{(0, \infty)}^{\oplus} V_a \otimes I_a d\mu(a) \oplus V_\infty \otimes I_\infty \oplus V_{-\infty} \otimes I_{-\infty} \oplus \int_{\mathbb{R}}^{\oplus} v_\lambda \otimes I_\lambda d\nu(\lambda),$$

where $V_a, V_\infty, V_{-\infty}$ are scalar operators.

We will prove in the next section that this mapping is in fact a bijection.

5. Unbounded generator of $C^*(\mathcal{G})$

Let U_t be a strongly continuous semigroup of partial isometries on H . Let us recall that for such a semi-group the (infinitesimal) generator A is defined by

$$A\varphi = \lim_{t \rightarrow 0} \frac{1}{t} (U_t - I)\varphi$$

for all φ whenever this limit exists. It is known that A is a closed operator with domain $\mathcal{D}(A)$ which is dense in H and A determines the semigroup uniquely (see [11]).

In this section we construct an unbounded element A which generates $C^*(\mathcal{G})$ as affiliated element (in the sense of [15,16]) and such that for each non-degenerate representation π of $C^*(\mathcal{G})$, $\pi(A)$ is the generator of the corresponding semigroup $\{U_t\}_{t \geq 0}$ of partial isometries.

First we recall some important definitions and results from [15,16]. Let \mathcal{A} be a C^* -algebra and T be a linear mapping acting on \mathcal{A} defined on a dense linear domain $D(T)$. The adjoint mapping is defined by the equivalence $(x, y \in \mathcal{A})$:

$$x \in D(T^*) \quad \text{and} \quad y = T^*x \quad \Leftrightarrow \quad \text{for any } a \in D(T) \quad (Ta)^*x = a^*y.$$

Let $B(\mathcal{A})$ be the algebra of bounded linear mappings acting on \mathcal{A} . Then $b \in B(\mathcal{A})$ is the Hermitean adjoint of $a \in B(\mathcal{A})$ and write $b = a^*$ if

$$y^*(ax) = (by)^*x$$

for any $x, y \in \mathcal{A}$. The set

$$M(\mathcal{A}) = \{a \in B(\mathcal{A}) : a^* \in B(\mathcal{A})\}$$

endowed with natural algebraic operations and the sup-norm is the multiplier algebra of \mathcal{A} . Assuming that elements of \mathcal{A} act on \mathcal{A} by left multiplication we have the embedding $\mathcal{A} \hookrightarrow M(\mathcal{A})$.

Let T be a linear mapping acting on \mathcal{A} having a dense domain $D(T)$. We say that T is *affiliated* with \mathcal{A} and write $T\eta\mathcal{A}$ if there exists $z \in M(\mathcal{A})$ such that $\|z\| \leq 1$, $D(T) = \sqrt{I - z^*z}\mathcal{A}$ and $T\sqrt{I - z^*z}a = za$ for all $a \in \mathcal{A}$. It is known that z is determined by T and one calls it z -transform of T and denotes it by z_T . One has that $T\eta\mathcal{A}$ implies $T^*\eta\mathcal{A}$ and $z_{T^*} = z_T^*$. If \mathcal{A} is a C^* -algebra of operators on H and T is a closed densely defined operator T on H then

$$z_T = T(I + T^*T)^{-1}.$$

The affiliation relation in the theory of C^* -algebras was introduced by S. Baaj and P. Jungl in [2]. This new reference will be under number 2, see References. Let $\text{Rep}(\mathcal{A}, H)$ denote the set of all non-degenerate representations of \mathcal{A} in a Hilbert space H , i.e. all $*$ -algebra homomorphisms $\pi : \mathcal{A} \rightarrow B(H)$ such that $\pi(\mathcal{A})H$ is dense in H . Each such $\pi \in \text{Rep} \mathcal{A}$ admits the unique extension to a unital $*$ -algebra homomorphism $\pi : M(\mathcal{A}) \rightarrow B(H)$. If $T\eta\mathcal{A}$ and $\pi \in \text{Rep}(\mathcal{A}, H)$ then there exists a uniquely defined closed operator S acting on H such that $\pi(z_T) = z_S$. We write $\pi(T) = S$.

For a C^* -algebra \mathcal{B} of operators on H we set

$$\text{Mor}(\mathcal{A}, \mathcal{B}) = \{\pi \in \text{Rep}(\mathcal{A}, H) : \overline{\pi(\mathcal{A})\mathcal{B}} = \mathcal{B}\}.$$

Let T_1, \dots, T_n be elements affiliated with \mathcal{A} . One says that \mathcal{A} is generated by T_1, \dots, T_n if for any Hilbert space H , any C^* -algebra \mathcal{B} of operators in $B(H)$ and any $\pi \in \text{Rep}(\mathcal{A}, H)$ one has

$$\pi(T_i)\eta\mathcal{B} \quad \text{for any } i = 1, \dots, n \quad \Rightarrow \quad \pi \in \text{Mor}(\mathcal{A}, \mathcal{B}).$$

In order to find a generator of $C^*(\mathcal{G})$ we will need to prove several lemmas bellow.

For the representation W_a of $C^*(\mathcal{G})$ we denote by $W_a(A)$ the infinitesimal operator of the corresponding semigroup. By [12], we have that $\mathcal{D}(W_a(A))$ consists of all absolutely continuous functions $f \in L_2([0, a])$, if $a \neq \infty$, and $f \in L_2([0, \infty))$, if $a = \infty$, satisfying $f(0) = 0$, and $(W_a(A)f)(x) = -f'(x)$, $f \in \mathcal{D}(W_a(A))$.

Let

$$h(x, t, a) = \begin{cases} -\chi_{[0,x]}(t) \sinh t + \frac{\sinh x}{\cosh a} \cosh(x-t-a), & a < \infty, \\ -\chi_{[0,x]}(t) \sinh t + \sinh x e^{t-x}, & a = \infty, x \neq \infty, \\ e^{-|t|}/2, & a = \infty, x = \infty, \end{cases}$$

$$\tilde{h}(x, t, a) = \begin{cases} \chi_{[0,x]}(t) \cosh t - \frac{\cosh(x-t-a) \cosh x}{\cosh a}, & a < \infty, \\ \chi_{[0,x]}(t) \cosh t - \cosh x e^{t-x}, & a = \infty, x \neq \infty, \\ -e^{-|t|}/2, & a = \infty, x = \infty, \end{cases}$$

$$k(x, t, a) = \begin{cases} -\chi_{[0,x]}(t) \sinh t - \frac{\cosh x}{\cosh a} \sinh(x-t-a), & a < \infty, \\ -\chi_{[0,x]}(t) \sinh t + \cosh x e^{t-x}, & a = \infty, x \neq \infty, \\ e^{-|t|}/2, & a = \infty, x = \infty, \end{cases}$$

$$\tilde{k}(x, t, a) = \begin{cases} -\chi_{[0,x]}(t) \cosh t - \frac{\sinh(x-t-a) \sinh x}{\cosh a}, & a < \infty, \\ -\chi_{[0,x]}(t) \cosh t + \sinh x e^{t-x}, & a = \infty, x \neq \infty, \\ -e^{-|t|}/2, & a = \infty, x = \infty. \end{cases}$$

Lemma 1. *The elements $h, \tilde{h}, k, \tilde{k}$ belong to $C^*(\mathcal{G})$. Moreover,*

$$W_a(h) = (1 + W_a(A)^* W_a(A))^{-1}, \quad W_a(\tilde{h}) = W_a(A) (1 + W_a(A)^* W_a(A))^{-1},$$

$$W_a(k) = (1 + W_a(A) W_a(A)^*)^{-1}, \quad W_a(\tilde{k}) = W_a(A)^* (1 + W_a(A) W_a(A)^*)^{-1},$$

for any $a \in (0, \infty]$.

Proof. To show the first statement it suffices to see that $\int_{x-a}^x |h(x, t, a)| dt$, $\int_{x-a}^x |\tilde{h}(x-t, -t, a)| dt$, $\int_{x-a}^x |\tilde{h}(x, t, a)| dt$ and $\int_{x-a}^x |\tilde{h}(x-t, -t, a)| dt$ are uniformly bounded on $\{(x, a): x \in [0, a], a > 0\}$. We have that

$$\begin{aligned} \int_{x-a}^x |h(x, t, a)| dt &= \frac{2 \cosh(a-x) \sinh^2(x/2)}{\cosh a} + \frac{\sinh x \sinh(a-x)}{\cosh a} \\ &\leq \frac{\cosh a - 1}{\cosh a} + \frac{\cosh a - 1}{2 \cosh a} \leq \frac{3}{2}. \end{aligned}$$

Similar calculations give us the uniform boundedness of $\int_{x-a}^x |\tilde{h}(x, t, a)| dt$.

Then $(W_a(A)^* W_a(A) f)(x) = -f''(x)$ with the domain consisting of $f \in L_2([0, a])$ such that f and f' are absolutely continuous, $f', f'' \in L_2([0, a])$ satisfy the boundary conditions $f(0) = 0$ and $f'(a) = 0$ if $0 \leq a < \infty$ and $f(0) = 0$ if $a = +\infty$.

Let $f, g \in L^2([0, a])$, $f \in \mathcal{D}(W_a(A)^* W_a(A))$ and $((1 + W_a(A)^* W_a(A)) f)(x) = -g(x)$. Then $f''(x) - f(x) = g(x)$ with general solution given by

$$f(x) = (e^x - e^{-x})h_1 + (e^x + e^{-x})h_2 + \frac{1}{2} \int_0^x e^{-(t-x)} g(t) dt - \frac{1}{2} \int_0^x e^{t-x} g(t) dt, \quad (7)$$

where h_1, h_2 are constants depending on g . Let first $0 < a < +\infty$. Since $f \in \mathcal{D}(W_a(A)^* W_a(A))$, $f(0) = 0$ and $f'(a) = 0$ which imply $h_2 = 0$ and $h_1 = -\frac{1}{2 \cosh a} \int_0^a \cosh(t-a)g(t) dt$.

Thus

$$\begin{aligned} & (((1 + W_a(A)^* W_a(A))^{-1})g)(x) \\ &= \frac{\sinh x}{\cosh a} \int_0^a \cosh(t-a)g(t) dt - \int_0^x \sinh(x-t)g(t) dt \\ &= \frac{\sinh x}{\cosh a} \int_{x-a}^x \cosh(x-t-a)g(x-t) dt - \int_{x-a}^x \chi_{[0,x]}(t) \sinh t g(x-t) dt \\ &= \int_{\mathbb{R}} h(x, t, a) (U_t^a g)(x) dt = (W_a(h)g)(x), \end{aligned}$$

where

$$h(x, t, a) = -\chi_{[0,x]}(t) \sinh t + \frac{\sinh x}{\cosh a} \cosh(x-t-a).$$

If $a = \infty$, one chooses $h_2 = 0$ and $h_1 = -\frac{1}{2} \int_0^\infty e^{-t} g(t) dt$ and obtains the equality $(1 + W_a(A)^* W_a(A))^{-1} = W_a(h)$. Next we calculate $W_a(A)(1 + W_a(A)^* W_a(A))^{-1}$. For $0 < a < +\infty$,

$$\begin{aligned} & (W_a(A)(1 + W_a(A)^* W_a(A))^{-1}g)(x) \\ &= -\left(\frac{\sinh x}{\cosh a} \int_0^a \cosh(t-a)g(t) dt - \int_0^x \sinh(x-t)g(t) dt \right)' \\ &= -\frac{\cosh x}{\cosh a} \int_0^a \cosh(t-a)g(t) dt + \int_0^x \cosh(x-t)g(t) dt \\ &= \int_{\mathbb{R}} \tilde{h}(x, t, a) (U_t^a g)(x) dt = (W_a(\tilde{h})g)(x). \end{aligned}$$

In a similar way one checks the case $a = \infty$. To show the statement about k , and \tilde{k} take $f, g \in L^2([0, a])$, $f \in \mathcal{D}(W_a(A)W_a(A)^*)$ and $((1 + W_a(A)W_a(A)^*)f)(x) = -g(x)$. Then $f''(x) - f(x) = g(x)$ as in the previous case with general solution given by (7). If $0 < a < +\infty$ we have for $f \in \mathcal{D}(W_a(A)W_a(A)^*)$ that $f(a) = 0$ and $f'(0) = 0$ implying

$$h_1 = 0 \quad \text{and} \quad h_2 = \frac{1}{2 \cosh a} \int_0^a \sinh(t-a)g(t) dt.$$

If $a = +\infty$ one has that any $f \in \mathcal{D}(W_a(A)W_a(A)^*)$ satisfies $f'(0) = 0$. To get also a solution $f \in L_2([0, \infty))$ one chooses $h_1 = 0$ and $h_2 = -\frac{1}{2} \int_0^\infty e^{-t} g(t) dt$. Similar verification gives

$$((1 + W_a(A)W_a(A)^*)^{-1}g)(x) = \int_{\mathbb{R}} k(x, t, a)(U_t^a g)(x) dt = (W_a(k)g)(x)$$

and

$$(W_a(A)^*(1 + W_a(A)W_a(A)^*)^{-1}g)(x) = \int_{\mathbb{R}} \tilde{k}(x, t, a)(U_t^a g)(x) dt = (W_a(\tilde{k})g)(x). \quad \square$$

As $W_a(h) = (1 + W_a(A)^*W_a(A))^{-1}$ is a positive operator for each $a > 0$ and $\pi = \int_{a>0}^\oplus W_a da$ is a faithful representation of $C^*(\mathcal{G})$, h is a positive element of $C^*(\mathcal{G})$. Consider the linear subspace $U = \{h^{1/2}b, b \in C^*(\mathcal{G})\}$ (here $h^{1/2}$ is the square root of h in $C^*(\mathcal{G})$) and define an operator z from U to $C^*(\mathcal{G})$ by

$$zh^{1/2}b = \tilde{h}b.$$

Lemma 2. *The subspace U is dense in $C^*(\mathcal{G})$. The mapping z can be extended to an element $z \in M(C^*(\mathcal{G}))$ such that $\|z\| \leq 1$ and $\sqrt{1 - z^*z}C^*(\mathcal{G})$ is dense in $C^*(\mathcal{G})$.*

Proof. Assume that U is not dense in $C^*(\mathcal{G})$. Then the closure \bar{U} is a proper right ideal. By [6, 2.9.5] there exists an irreducible representation π on H and a cyclic vector $\varphi \in H$ such that $(\pi(h^{1/2}b)\varphi, \varphi) = 0$, $b \in C^*(\mathcal{G})$ implying $(\pi(b)\varphi, \pi(h^{1/2})\varphi) = 0$ and $\pi(h^{1/2})\varphi = 0$. By Theorem 5 any irreducible representation π is unitarily equivalent to one from the set $S = \{W_a, \omega_\phi, a \in \mathbb{R}^+, \phi \in \mathbb{R}\}$. If π is unitarily equivalent to W_a , ($a \in (0, \infty]$), then $\ker \pi(h^{1/2}) = \{0\}$ since $\ker W_a(h^{1/2}) = \ker(1 + W_a(A)^*W_a(A))^{-1} = \{0\}$. For $\pi = \omega_\phi$ we have $\pi(h) = 1/2$ and therefore $\ker \omega_\phi(h^{1/2})$ is also trivial. A contradiction. As

$$\begin{aligned} \|zh^{1/2}b\| &= \sup_{a>0} \|W_a(zh^{1/2}b)\| = \sup_{a\geq 0} \|W_a(\tilde{h}b)\| \\ &= \sup_{a>0} \|(W_a(A)(1 + W_a(A)^*W_a(A))^{-1/2})W_a(h^{1/2})W_a(b)\| \\ &\leq \sup_{a>0} \|W_a(h^{1/2}b)\| = \|h^{1/2}b\|, \end{aligned}$$

z can be extended to a bounded operator on $\bar{U} = C^*(\mathcal{G})$ with the norm $\|z\| \leq 1$. We have also that for $\pi = W_a$ on a Hilbert space H the equality $\pi(z)\pi(h)^{1/2}\pi(b)\varphi = \pi(\tilde{h})\pi(b)\varphi$ is equivalent to

$$W_a(z)(1 + W_a(A)^*W_a(A))^{-1/2}W_a(b)\varphi = W_a(A)(1 + W_a(A)^*W_a(A))^{-1}W_a(b)\varphi$$

giving us that $W_a(z)\psi = W_a(A)(1 + W_a(A)^*W_a(A))^{-1/2}\psi$ for any $\psi \in H$. Here we denote by π also the extension of a non-degenerate representation π of $C^*(\mathcal{G})$ to the space of bounded linear mappings on $C^*(\mathcal{G})$.

To prove that $z \in M(C^*(\mathcal{G}))$ we have to show that z^* is also bounded. Consider the set $\tilde{U} = \{k^{1/2}a: a \in C^*(\mathcal{G})\}$. In a similar way one shows that \tilde{U} is dense in $C^*(\mathcal{G})$. Define an operator

\tilde{z} on \tilde{U} by $\tilde{z}k^{1/2}a = \tilde{k}a$ for all $a \in C^*(\mathcal{G})$. As above one checks that \tilde{z} can be extended to a bounded mapping on $C^*(\mathcal{G})$ with the norm $\|\tilde{z}\| \leq 1$. One also has that $W_a(\tilde{z})\psi = W_a(A)^*(1 + W_a(A)W_a(A)^*)^{-1/2}\psi$ for any $\psi \in H$. Our aim now is to show that $z^* = \tilde{z}$. Take $y \in \tilde{U}$, i.e., $y = k^{1/2}a$, $a \in C^*(\mathcal{G})$, and $x \in U$, i.e., $x = h^{1/2}b$, $b \in C^*(\mathcal{G})$. We have to prove the equality $y^*(zx) = (\tilde{z}y)^*x$, i.e., $b^*k^{1/2}\tilde{h}a = b^*\tilde{k}^*h^{1/2}a$. For this it is enough to see that $k^{1/2}\tilde{h} = \tilde{k}^*h^{1/2}$. Applying the representation W_a to the equality we obtain

$$\begin{aligned} & (1 + W_a(A)W_a(A)^*)^{-1/2}W_a(A)(1 + W_a(A)^*W_a(A))^{-1} \\ &= ((W_a(A))^*(1 + W_a(A)W_a(A)^*)^{-1})^*(1 + W_a(A)^*W_a(A))^{-1/2}. \end{aligned} \quad (8)$$

As $z_T = T(1 + T^*T)^{-1/2}$, we arrive that (8) is equivalent to $z_{W_a(A)}^* = z_{W_a(A)}^*$ which clearly holds.

Direct calculation shows now also that $\pi(\sqrt{1 - z^*z}) = (1 + W_a(A)^*W_a(A))^{-1/2} = \pi(h^{1/2})$. Thus $\sqrt{1 - z^*z} = h^{1/2}$ giving the density of $\sqrt{1 - z^*z}C^*(\mathcal{G})$. \square

Define now a linear map A acting on $C^*(\mathcal{G})$ with domain $D(A) = \sqrt{1 - z^*z}C^*(\mathcal{G})$ as follows: $A\sqrt{1 - z^*z}b = zb$ for all $b \in C^*(\mathcal{G})$. A is affiliated with $C^*(\mathcal{G})$ by the construction and $z_A = z$.

Proposition 7. *The mapping $F: \text{Rep}_+(U, H) \rightarrow \text{Rep}(C^*(\mathcal{G}), H)$ is a bijection.*

Proof. A semigroup $\{U_t\}_{t \geq 0} \in \text{Rep}_+(U, H)$ is uniquely defined by its generator \mathcal{A} . It is easy to deduce from the definition of F that the z -transform $z_{\mathcal{A}}$ is equal to $\pi(z)$ where π is a unique extension of representation $F(\{U_t\})$ of the C^* -algebra $C^*(\mathcal{G})$ to its multiplier algebra $M(C^*(\mathcal{G}))$ and $z \in M(C^*(\mathcal{G}))$ is the z -transform of the affiliated element A defined above. Since the correspondence $\mathcal{A} \rightarrow z_{\mathcal{A}}$ is one-to-one we conclude that F is an injection. That F is a surjection follows from Theorem 6. \square

Theorem 8. *The linear mapping A generates $C^*(\mathcal{G})$ as affiliated element. If π is a representation of $C^*(\mathcal{G})$ on H , then $\pi(A)\varphi = A_\pi\varphi$, for all $\varphi \in \pi(D(A))H$, where A_π is the generator of the corresponding semigroup $\{U_t\}_{t \geq 0}$ of partial isometries.*

Proof. To prove that A generates the algebra, by [16, Theorem 3.3], it is enough to show that A separate representations of $C^*(\mathcal{G})$, i.e. if π_1, π_2 are different non-degenerate representations of $C^*(\mathcal{G})$ in a Hilbert space H then $\pi_1(A) \neq \pi_2(A)$, and that $(1 + A^*A)^{-1} \in C^*(\mathcal{G})$. The first claim follows from Proposition 7 and the fact that a semigroup is uniquely defined by its generator. The second follows from the equality $(1 + A^*A)^{-1} = 1 - z^*z = h$ obtained in Lemma 2.

For a representation π we have $\pi(A)\pi(\sqrt{1 - z^*z})\pi(b)\varphi = \pi(z)\varphi$. As $\pi(z) = z_{A_\pi} = A_\pi(1 + A_\pi^*A_\pi)^{-1/2}$ and $\pi(\sqrt{1 - z^*z}) = (1 + A_\pi^*A_\pi)^{-1/2}$ we have the required equality between $\pi(A)$ and A_π . \square

Acknowledgments

The work was partially written when the first and the second authors were visiting Chalmers University of Technology in Göteborg, Sweden. The research was partially supported by a grant from the Swedish Royal Academy of Sciences and STINT as a part of the program of cooperation with the former Soviet Union. The second author was also supported by the Swedish Institute. The third author was supported by the Swedish Research Council.

Appendix A

Let \mathcal{K} be the algebra of compact operators on $H = L_2([0, +\infty))$ and let $\mathcal{A} = C_0((0, +\infty], \mathcal{K})$. Denote by H_a the Hilbert space $L_2([0, a])$ which we consider as a subspace of H and let

$$\mathcal{B} = \{f \in \mathcal{A}: f(a)H_a \subset H_a, f(a)|_{H_a^\perp} = 0\}.$$

Proposition A.1.

$$\mathcal{B} \simeq \mathcal{A}.$$

Proof. Let us denote by \mathcal{K} also the algebra of all compact operators on $L_2([0, 1])$. Clearly, the algebra $\mathcal{C} = C_0((0, 1], \mathcal{K})$ is isomorphic to the algebra \mathcal{A} . Let for $0 < t \leq 1$, $V_t = L_2([0, t])$. Considering V_t as subspaces of $L_2([0, 1])$ we let

$$\mathcal{D} = \{g \in \mathcal{C}: g(t)V_t \subset V_t, g(t)|_{V_t^\perp} = 0\}.$$

Proving that \mathcal{B} is isomorphic to \mathcal{D} and then that \mathcal{D} is isomorphic to \mathcal{C} we get the statement of the proposition. Let $S: L_2([0, +\infty)) \rightarrow L_2([0, 1])$ be the isometric isomorphism given by $(S\xi)(t) = \frac{1}{1-t}\xi(\frac{t}{1-t})$. We have

$$SH_a = V_{\frac{a}{1+a}}, \quad a > 0. \quad (\text{A.1})$$

Define now a mapping $\varphi: \mathcal{B} \rightarrow \mathcal{C}$ by

$$\varphi(f)(a) = Sf\left(\frac{a}{1-a}\right)S^*, \quad a \in (0, 1].$$

By (A.1) $\varphi(f)(a)V_a \subset V_a$ and $\varphi(f)(a)|_{V_a^\perp} = 0$ and therefore $\varphi(\mathcal{B}) \subseteq \mathcal{D}$. One can easily see that φ is even a $*$ -isomorphism between \mathcal{B} and \mathcal{D} . Define $S_a: V_a \rightarrow L_2([0, 1])$ as follows

$$(S_a\xi)(x) = \sqrt{a}\xi(ax).$$

Then S_a is a unitary operator. For $f \in \mathcal{D}$ we set

$$\psi(f)(a) = S_af(a)S_a^*.$$

For each $a \in (0, 1]$, $\psi(f)(a) \in \mathcal{K}$. To prove that $\psi(f) \in \mathcal{C}$ we have to show that the mapping $a \mapsto \psi(f)(a)$ is continuous. We check this first for $f = T_{\eta, \zeta}$, where

$$\eta, \zeta \in \mathcal{X} := \{\xi \in C((0, 1], C[0, 1]): \xi_a \in V_a, a \in (0, 1]\} \quad \text{and} \\ (T_{\eta, \zeta}(a)\xi)(x) = (T_{\eta_a, \zeta_a}\xi)(x) = (\xi, \eta_a)\zeta_a(x).$$

We have

$$S_aT_{\eta_a, \zeta_a}S_a^*\xi = (S_a^*\xi, \eta_a)S_a\zeta_a = (\xi, S_a\eta_a)S_a\zeta_a$$

and for $a, b \in (0, 1]$

$$\begin{aligned} \|S_a T_{\eta_a, \zeta_a} S_a^* \xi - S_b T_{\eta_b, \zeta_b} S_b^* \xi\| &= \|(\xi, S_a \eta_a) S_a \zeta_a - (\xi, S_b \eta_b) S_b \zeta_b\| \\ &\leq \|(\xi, S_a \eta_a - S_b \eta_b) S_a \zeta_a\| + |(\xi, S_b \eta_b)| \cdot \|S_a \zeta_a - S_b \zeta_b\| \\ &\leq \|S_a \eta_a - S_b \eta_b\| \cdot \|\zeta_a\| \cdot \|\xi\| + \|S_a \zeta_a - S_b \zeta_b\| \cdot \|\eta_b\| \cdot \|\xi\| \end{aligned}$$

so that

$$\|S_a T_{\eta_a, \zeta_a} S_a^* - S_b T_{\eta_b, \zeta_b} S_b^*\| \leq \|S_a \eta_a - S_b \eta_b\| \cdot \|\zeta_a\| + \|S_a \zeta_a - S_b \zeta_b\| \cdot \|\eta_b\|.$$

Further

$$\|S_a \eta_a - S_b \eta_b\|^2 = \int_0^1 |\sqrt{a} \eta_a(ax) - \sqrt{b} \eta_b(bx)|^2 dx \rightarrow 0 \quad \text{as } a \rightarrow b$$

since $\eta(a, x) = \eta_a(ax)$ is continuous. Similarly, $\|S_a \zeta_a - S_b \zeta_b\| \rightarrow 0$ as $a \rightarrow b$. Moreover, $\|\zeta_a\|$ is bounded in a neighborhood of b giving $\|S_a T_{\eta_a, \zeta_a} S_a^* - S_b T_{\eta_b, \zeta_b} S_b^*\| \rightarrow 0$ as $a \rightarrow b$, i.e. $\psi(T_{\eta, \zeta})$ is continuous.

Next we prove that the linear span, \mathcal{M} , of elements $T_{\eta, \zeta}$ is dense in \mathcal{D} . Since \mathcal{D} is postliminal, it is enough to prove that \mathcal{M} is rich [6, 11.1]. Note first that \mathcal{D} is a subalgebra of \mathcal{C} and since any irreducible representation of \mathcal{C} is unitary equivalent to $\pi_t(f) = f(t)$, $f \in \mathcal{C}$ one can easily obtain that any irreducible representation of \mathcal{D} is given by $\pi'_t: f \mapsto f(t)|_{V_t}$ up to unitary equivalence. For any fixed $t \in (0, 1]$ we have $\pi'_t(T_{\eta, \zeta}) = T_{\eta_t, \zeta_t}$. Moreover, for any $\eta', \zeta' \in C([0, t]) \subset L_2([0, t])$ there exist $\eta, \zeta \in \mathcal{X}$ such that $\eta_t = \eta'$ and $\zeta_t = \zeta'$. For example, one can take $\eta_t = S_t^* \eta'$, $\zeta_t = S_t^* \zeta'$. Hence $\pi_t(\mathcal{M})$ is dense in the algebra of compact operators on $L_2([0, t])$ ensuring that π'_t is irreducible as representation of \mathcal{M} . Next we claim that for any pair of non-equivalent irreducible representations π_1, π_2 of \mathcal{D} (up to a unitary equivalence we have that $\pi_1 = \pi'_t, \pi_2 = \pi'_s$ for distinct $t, s \in (0, 1]$) the restrictions $\pi_1|_{\mathcal{M}}, \pi_2|_{\mathcal{M}}$ are also non-equivalent. For this it is enough to see that there is no unitary operator $U: L_2([0, s]) \rightarrow L_2([0, t])$ such that $U T_{\eta_s, \zeta_s} U^* = T_{\eta_t, \zeta_t}$, i.e. $U \zeta_s = \zeta_t$, $U \eta_s = \eta_t$, for any $\eta, \zeta \in \mathcal{X}$. Take $\eta \in C([0, 1])$ such that $\text{supp } \eta = [0, 1]$. Assume $t < s$ and take arbitrary $b \in (t, s)$. Set

$$\eta_x = S_x^* \eta \quad \text{and} \quad \zeta_x = \begin{cases} S_x^* \eta, & x < b, \\ S_b^* \eta, & x \geq b. \end{cases}$$

It is easy to check that $\eta, \zeta \in \mathcal{X}$. Clearly $\eta_t = \zeta_t$ but since $\text{supp } \eta_s = [0, s] \neq \text{supp } \zeta_s = [0, b]$ we obtain $\eta_s \neq \zeta_s$ which gives the claim. Therefore \mathcal{M} is rich and consequently dense in \mathcal{D} giving $\psi(\mathcal{D}) \subset \mathcal{C}$.

Using similar argument one can obtain that $\psi(\mathcal{D})$ is a rich subalgebra of \mathcal{C} and therefore is equal to \mathcal{C} . Hence ψ is surjective. It is also obvious that ψ is injective and therefore ψ is the required $*$ -isomorphism. \square

References

- [1] W. Arveson, Continuous analogues of Fock space. II. The spectral C^* -algebra, *J. Funct. Anal.* 90 (1) (1990) 138–205.
- [2] S. Baaj, P. Julg, Théorie bivariante de Kasparov et opérateurs non bornés dans les C^* -modules hilbertiens (Bivariant Kasparov theory and unbounded operators on Hilbert C^* -modules), *C. R. Acad. Sci. Paris Sér. I Math.* 296 (21) (1983) 875–878 (in French).
- [3] L.A. Coburn, C^* -algebras generated by semi-groups of isometries, in: *Proc. Internat. Conf. Hilbert Space Operators and Operator Algebras*, Tihany, 1970, in: *Colloq. Math. Soc. Janos Bolyai*, vol. 5, North-Holland, Amsterdam, 1972, pp. 99–105.
- [4] L.A. Coburn, R.G. Douglas, C^* -algebras of operators on a half-space. I, *Inst. Hautes Études Sci. Publ. Math.* 40 (1971) 59–67.
- [5] L.A. Coburn, R.G. Douglas, D.G. Schaeffer, I.M. Singer, C^* -algebras of operators on a half-space. II. Index theory, *Inst. Hautes Études Sci. Publ. Math.* 40 (1971) 69–79.
- [6] J. Dixmier, C^* -Algebras, North-Holland, Amsterdam, 1977 (English translation).
- [7] R.G. Douglas, On the C^* -algebra of a one-parameter semigroup of isometries, *Acta Math.* 128 (3–4) (1972) 143–151.
- [8] M.R. Embry, A.L. Lambert, L.J. Wallen, A simplified treatment of the structure of semigroups of partial isometries, *Michigan Math. J.* 22 (1975) 175–179.
- [9] R. Exel, M. Laca, J. Quigg, Partial dynamical systems and C^* -algebras generated by partial isometries, *J. Operator Theory* 47 (1) (2002) 169–186.
- [10] P.S. Muhly, J.N. Renault, C^* -algebras of multivariable Wiener–Hopf operators, *Trans. Amer. Math. Soc.* 274 (1982) 1–44.
- [11] B. Sz. Nagy, C. Foias, *Harmonic Analysis of Operators on Hilbert Space*, North-Holland, Amsterdam, 1970, xiii+389 pp.; translated from French and revised.
- [12] V. Ostrovskiy, Centered one-parameter semigroups, *Methods Funct. Anal. Topology* 10 (2) (2004) 32–42.
- [13] A. Paterson, *Groupoids, Inverse Semigroups and Their Operator Algebras*, Birkhäuser, Boston, MA, 1999, xvi+274 pp.
- [14] J. Renault, *A Groupoid Approach to C^* -Algebras*, *Lecture Notes in Math.*, vol. 793, Springer-Verlag, Berlin, 1980, ii+160 pp.
- [15] S.L. Woronowicz, Unbounded elements affiliated with C^* -algebras and noncompact quantum groups, *Comm. Math. Phys.* 136 (2) (1991) 399–432.
- [16] S.L. Woronowicz, C^* -algebras generated by unbounded elements, *Rev. Math. Phys.* 7 (3) (1995) 481–521.